

Authors: *Dr. Eric P. Berg*
University of Missouri

Reviewer: *Dr. Scott J. Eilert*
Excel Corporation

Critical Points Affecting Fresh Pork Quality within the Packing Plant

Introduction

Postmortem metabolism of porcine muscle is more rapid than beef or lamb (Marsh *et al.* 1972). Metabolism of intramuscular glycogen plays the primary role in the conversion of muscle to meat and the expression of different quality attributes of fresh pork. The two most common quality concerns with fresh pork are pale, soft, and exudative (PSE) and dark, firm, and dry (DFD) lean, with PSE being much more economically important than DFD. Both are a result of *postmortem* glycogen metabolism. The level and extent of *postmortem* pH decline is glycogen dependent as anaerobic conversion of glycogen to lactic acid results in the achievement of an acceptable (or unacceptable) meat pH. The PSE condition occurs when intramuscular lactic acid (localized lactidosis) accumulates very rapidly (< 1h) while the carcass temperature is still high. The initiation of glycolysis could be attributed to many things: **1)** genetic predisposition (porcine stress syndrome); **2)** elevated metabolism or if a pig is prone to a high "excitability" (Grandin, 1994); **3)** pre-slaughter stress; and (or) **4)** combinations of all of these. Activation of the glycolytic system just prior to exsanguination leads to the production of heat which will elevate the pig's body temperature. Pearson (1987) wrote that the ultimate condition of pork muscle is influenced by skeletal muscle pH drop as a function of time, *in vivo* temperature patterns, carcass chilling rate, and the conditions at the onset of rigor mortis. The rate of *postmortem* pH decline is approximately three times faster in carcasses that ultimately produce PSE meat, regardless of presence or absence of the stress gene (Mitchell and Heffron, 1982). This rapid pH drop while the carcass temperature is above 37°C leads to denaturation of myofibrillar proteins that bind water (Penny, 1969).

Grandin (1994) reported that ultimate pork quality was a responsibility given 50% to the producer and 50% to the packer. The producer is responsible for selection of swine genotypes that possess acceptable pork quality traits and must provide environmental conditions to optimize *composition* (growth) and *quality* of the final food product. The producer must guarantee proper selection, care, and handling to the point of delivery to the packing plant. The packer accepts the responsibility of optimizing pre- and post-slaughter conditions to ensure optimal meat quality. The purpose of this review is to identify the critical points affecting fresh pork quality at the slaughter and processing plant with a brief discussion regarding pig handling at the production site. This review will provide scientific documentation regarding conditions which can be controlled within the packing plant in an effort to optimize fresh pork quality.

Production Site

Acceptable pork quality is the responsibility of both the packer and producer (Grandin, 1994) therefore, a brief discussion regarding conditions at the production site is in order. Appropriate genetic selection is but one attribute that can be controlled by the swine producer. Adequate care and treatment must be a standard production practice. Good management includes development of an efficient and economic feed ration, proper record keeping (sire and dam summaries), and appropriate housing conditions. Grandin (1989) suggested methods for "environmental enrichment" of feeder pigs for growth in a less stressful environment. Pigs that are accustomed to people and sounds are less likely to produce quality problems associated with pre-slaughter handling. Hogs become more accustomed to sudden noises if they have been raised in an environment

where a radio has been playing or have had frequent encounters with people (such as opening and closing pen gates; walking in pens). Excitable pigs (Grandin 1998a) and pigs that have had little human contact or poor "environmental enrichment" will experience more stress during loading and transport. Hunter *et al.* (1997) found that the rearing, finishing, and loading environment will significantly affect the stress encountered by transport and lairage. Pigs which are easy to move are less likely to be treated harshly during transport and lairage.

Pre-transport handling at the production site.

Pigs react to a stress with an inborn fear response. High levels of fear may result in a physiological stress response that may adversely affect welfare, productivity, and ultimately meat quality (Hemsworth, 1993). Transport is stressful for pigs (Mickwitz,

1982; van Putten, 1982; McGlone *et al.*, 1993; Lambooi and van Putten, 1993; Martoccia, *et al.* 1995), therefore, care should be taken to minimize handling stress prior to transport. Kilgour and Dalton (1984) suggest holding market pigs together in an unfamiliar pen the night before transport and moving them early in the morning. Pigs will load more easily into a trailer if they are provided secure footing and the trailer is level with the loading plane. Market weight pigs are more difficult to drive on concrete slabs, plastic, or metal floors because of unstable footing (Warris *et al.*, 1983; Grandin, 1994). Pigs will load more easily through a dual ramp and chute design constructed to facilitate side by side loading. A distressed pig will try to maintain contact with its pen-mate if not physically, then at least visually (van Putten, 1982). The outer sides of the loading ramp

continued on page 2

continued from page 1

should be solid with an inner middle partition panel that is “see-through” (such as a wire panel). The see-through middle panel will promote the pigs’ following behavior while the solid outer sides restrict the view of distractions outside of the chute (Grandin, 1998b; Grandin, 1998c). Because sight restriction lowers stress levels, a solid portable (hand-held) panel is a very effective means of driving market

hogs. Electric prods should be avoided and used only in cases of emergency (Grandin, 1989; Guise and Penny, 1989a; Kilgour and Dalton, 1984). These same guidelines are effective for unloading pigs upon arrival at the packing plant.

Pigs have a tendency to move from dimly to brightly lit areas. A spotlight directed into a ramp or trailer will facilitate entry as long as it illuminates the interior of the ramp or trailer and does not shine directly in the pigs eyes (Grandin, 1989).

Nanni Costa *et al* (1996) reported that use of a loading ramp that has a minimal slope had no negative consequences on muscle quality. Tarrant (1989) recommends these options for loading of pigs (listed in order of importance): **1)** a loading platform that is level with the truck deck; **2)** a hydraulic tailgate lift on the truck; **3)** a fixed, permanent loading ramp with an incline slope of 20° (pigs have an aversion to downward slopes); **4)** a portable loading ramp with an incline of 27°.

Transport

Martocchia *et al.* (1995) concluded that pig transport was the most influential pre-slaughter factor because it affects both meat quantity and quality. Grandin (1998d) provides several recommendations to reduce transport stress in market swine. Trailers should be maintained and kept in good repair. Trucks and trailers should be cleaned after each load to prevent slipping and skin blemishes (carcass bruising). Overcrowding will result in increased body temperature (Guise and Warris, 1989), death loss, and PSE carcasses.

Studies of transportation stress are difficult to analyze because of the cumulative contribution of each component associated with transport (Stephens and Perry, 1990). Changes in velocity, vibration, handling by unfamiliar persons, mixing with unfamiliar hogs, and establishment of a new social group all contribute to the total stress of transport. The factor (or combination of factors) contributing the highest amount of total stress may not be readily evaluated. In general, transportation conditions will affect post-mortem meat quality by provoking stress or animal fatigue (Lambooj and van Putten, 1993).

Stocking density, weather, and temperature during transport

The person transporting pigs will most often determine the stocking density of the truck or trailer. The number of pigs per load is largely a function influenced by economics. Market pigs will experience heat stress, fatigue, lower meat quality, and a higher mortality if too little space is allowed. Too much space is not cost effective and the extra space provides more room for injury as pigs can be thrown about during sudden stops and cornering (Tarrant, 1989). The effect of stocking density on ultimate meat quality is greatly

influenced by breed and ambient temperature. Tarrant (1989) provides a review of different stocking densities used for road transport of pigs to slaughter. The average, minimum recommended space allowance of 0.35 m²/100 kg pig was derived from the review of 10 transport studies. The space allowance per pig must be increased by 10% in hot weather and when the route to slaughter is through areas of heavy traffic or urban areas where ventilation may be reduced as a result of slowed or stopped transport velocity.

Trucks or trailers should be lined with wet sand or shavings to keep pigs cool when temperatures exceed 15° C (60° F). When temperatures are above 27° C (80° F) pigs should be sprinkled with water prior to loading. Never bed pigs with straw during hot weather. High temperature and humidity are extremely dangerous for pigs because of their inefficient means of heat respiration. In fact, Williams (1985; as cited by Tarrant, 1993) found that pig transport death rate plotted against temperature began a steep rise when temperatures exceeded 18°C and became almost vertical above 25°C.

Grandin (1998d) recommends pigs be scheduled for delivery to the packing plant at night or early morning when the Livestock Weather Safety Index is in the Danger and Emergency zone. If this is not possible, trucks should be kept moving to facilitate air flow. During hot weather, pigs should be unloaded as soon as they reach the plant. In cold weather at temperatures below 15° C (60° F), trucks and trailers should be bedded with straw or deep, dry shavings to keep pigs warm. Aluminum transport trailers should have approximately half the air flow holes closed with plywood if temperatures fall below minus 12° C (10° F). Eilert (1997) stated that from a physiologic standpoint, if a hog gets exces-

sively cold on a truck, this will not affect pork quality. However, if the pig experiences cold to the degree of frost bite or wind chill burn, the respective pork carcass will require more time on the kill floor for extra trimming (at an elevated temperature) which is detrimental to pork quality. Grandin (1994) advises that it is very important to take weather into account when testing new methods for reducing PSE and (or) bloodsplash at the packing plant. The incidence of PSE carcasses may easily double during the first four hot days of spring.

Length of Transit

Transit duration has a variable affect on pork quality. Martocchia *et al.* (1995) found that pigs transported 650 km had a significantly higher pH₄₅ (45 min *post-mortem* pH), pH_{ult} (ultimate pH), and a darker, redder meat than pigs transported 180 km. These authors are quick to conclude that, although these differences were significant, they were well within the range considered normal. Warris *et al.* (1990) found transportation times of 1 to 4 h had no significant affect on pH_{ult}. A large amount of the stress of transport occurs at loading and unloading. Bradshaw *et al.* (1996) reported that concentrations of the stress hormones cortisol and beta-endorphin were substantially elevated immediately after loading. Shorter transport time may be more detrimental than longer ones if the driving, stocking density, and ventilation are adequate (Tarrant, 1989). Stephens and Perry (1990) suggest that pigs may acclimate to the stress of transport. The authors found the largest increase in heart rate occurred immediately on starting the engine (to generate vibration and noise) and then gradually

continued on page 3

continued from page 2

declined as the test progressed. Grandin (1994) observed that pigs hauled very short distances for under 30 minutes are often more stubborn and difficult to drive at the plant than pigs experiencing a longer transit time and *short duration* stress may lead to a higher incidence of PSE. Pigs transported a longer distance may be more likely to develop DFD meat as a result of *long duration* stress and depletion of intramuscular glycogen stores.

Death loss due to transport was lowest for short transit duration (10 - 25 min) while the greatest death occurred during 45 - 80 minutes of transport (Honkavaara, 1989a) suggesting that stress sensitive pigs die in a relatively short period of time (< 45 min) as a result of transport. Honkavaara (1989a) found maximum mortality occurred in warm weather (14 to 24°C) while the least mortality occurred in cool weather (around 8°C).

Lairage

Martocchia *et al.* (1995) concluded that transport distance itself did not seem to determine severe levels of stress in pigs and found that transport stress may be offset by the amount of time pigs spend in lairage before stunning. It is generally accepted that time spent in lairage allows a stressed animal time to recover from loading, transport, and unloading stressors. Pigs should be rested (transport recovery period) 2 to 4 hours before entering the stunning chute (Grandin, 1994). Once unloaded into lairage, pigs will rapidly begin establishing a dominance hierarchy. Messe and Ewbank (1973) found that the average aggression rate was 2 incidents per pig per hour in groups of familiar pigs. Pigs in unfamiliar groups have higher frequency, higher intensity, more spontaneous, and less resource-related aggression. The average aggression rate can be as much as 12 incidents per pig per hour. Baxter (1985) reported that 10 groups of 6 previously unacquainted pigs gave an average of 138 aggressive interactions per group in the first hour after introduction (23 aggressive interactions per pig per hour). Pigs slaughtered during the initial hours of lairage, during this period of aggressive behavior, experience both

physical and physiological stress (Grandin, 1997) which will result in increased metabolic activity, reduced intramuscular pH (Enfalt *et al.*, 1993), and elevated body temperature. On average, the hierarchy is established within 2 hours and pigs begin to rest. Grandin (1994) suggests that after 4 h in lairage pigs begin to "wake up" and return to the aggressive behavior that may ultimately affect meat quality. Honkavaara (1989b) concluded that the high occurrence of PSE carcasses in his study may have been largely due to the stressful treatment of pigs being woken up and driven to the stunner after being held in lairage overnight. De Smet *et al.* (1996) found a lairage time of 2 to 6 h, compared with immediate slaughter, improved meat quality resulting in a lower percentage drip loss, lower Hunter L-values, and higher pH₄₅.

Warris (1987) provides an excellent review of the effect of time and conditions of transport and lairage on pig meat quality and contends that under modern conditions, holding pigs in lairage may have negligible or no influence on meat quality. Prolonged holding may lead to a reduction in meat quality if pigs have not been subjected to stressful handling prior to or during transport or if they are of a breed particularly resistant to stress. Warris *et al.* (1996) found longer lairage time reduced the incidence of PSE meat but increased the prevalence of DFD meat. Longer lairage produced progressively more skin blemishes as a result of fighting among unfamiliar pigs and an overnight lairage reduced carcass yield and backfat thickness. Warris (1987) summarizes Elliot and Patton (1968) who concluded that if pigs were not stressed on arrival at the plant, resting in lairage would not improve meat quality. However, if pigs experienced stressful transport, then a lairage of at least 2 h was suggested for groups of mixed animals. Warris *et al.* (1996) found no evidence that leaner, more stress-susceptible pigs reacted differently to length of time in lairage than fatter, more stress-resistant animals.

Water sprays

Schutte *et al.* (1996) found that showering pigs with water after transport possesses three distinct advantages: **1**) it cools the pigs, reducing the strain on the cardiovascular system; **2**) calms the pigs, reducing aggressive behavior in lairage; and **3**) cleans the pigs, reducing contamination in the slaughter line. Pigs in lairage should

have *ad libitum* availability of drinking water and (ideally) cool showers during hot periods of the year (Grandin, 1994; Weeding *et al.*, 1993). These precautions will lower muscle temperature prior to exsanguination allowing body temperature to be more close to "normal" during early *postmortem* metabolism conversion of muscle to meat (Klont and Lambooy, 1995). Offer (1991) found that a reduction in body temperature of 2°C prior to exsanguination would reduce the initial rate of denaturation of myosin by 37% resulting in a corresponding reduction in drip loss. An intermittent shower regime may provide the greatest cooling effect through evaporative cooling on the pigs skin surface (Weeding *et al.*, 1993). Pigs have no sweat glands in their skin and use water and wallowing as a means of keeping cool in hot climates (Kilgore and Dalton, 1984; Lambooy and van Putten, 1993). Weeding *et al.* (1993) observed that pigs sprayed with water during lairage exhibited less stereotypic stressful behavior. Water sprayed pigs were involved with rooting about the pen floor, chewing and licking pen walls, and walking about the pen. These results would suggest that water sprayed pigs experience better welfare which should facilitate greater ease of handling upon entrance into the stunning chute.

Honkavaara (1989b) suggested that optimum lairage temperature, humidity, and time were 15-18°C, 59-65%, and 3-5 h, respectively. These levels resulted in lower muscle lactate levels and decreased occurrence of PSE meat.

Feed Restriction.

It is generally accepted that a period of fasting before slaughter will reduce the amount of total carbohydrate available for *postmortem* conversion of glycogen to lactic acid (De Smet *et al.*, 1996). From an animal welfare standpoint, it may appear that the stress of transport combined with the stress of feed restriction may produce deleterious affect on the well being of pigs being transported to market. This is not the case. Becker *et al.* (1989) reported that the combination of feed restriction and transport stress did not present an additive demand on the pigs biological systems. In fact, Stephens and Perry (1990) found that a pigs aversion to transport was increased if they had eaten a large meal before transport and Tarrant (1993) reported that

continued on page 4

continued from page 3

death losses are greater when pigs are fed on the day of transport. Eilert (1997) suggested that producers factor in approximate transit time and holding time at the plant to determine length of feed withdrawal. Jones *et al.* (1985) reported overnight fasting at the plant resulted in increased ultimate meat pH, meat color intensity, and water holding capacity. Eikelenboom *et al.* (1991) found that hogs held off feed (but

not water) for a minimum of 12 h before slaughter exhibit fewer PSE carcasses. Murray *et al.* (1989) reported that a 24 or 48 h fast significantly reduced the frequency of PSE in stress susceptible (porcine stress syndrome) pigs. The advantage of feed restriction is not unanimous in the literature. De Smet *et al.* (1996) found overnight feed withdrawal to have no effect on meat quality for all PSE related traits. Becker *et al.* (1989) found no relationship between pigs fasted up to 72 hs

and ultimate pork quality but did note that meat tenderness was enhanced by transport and fasting.

Feed restriction of market pigs has a more practical application even if one discounts the effect on pork quality. Eilert (1997) suggests that a 16 to 24 h feed withdrawal period is consistently advantageous to the packer because of the greater ease of carcass evisceration, reduced waste handling at the plant, and reduced incidence of broken viscera on the slaughter floor.

Stunning

Movement of pigs to stunning

Grandin (1994) maintains that selection for lean, heavy muscled hogs has inadvertently resulted in increased problems with extremely excitable hogs that "are almost impossible to handle in a calm, gentle manner." As pig excitability has increased, problems with single file stunning chutes have worsened. It is natural behavior for cattle to move well through single-file chutes because cattle instinctively move from pasture to pasture via narrow, single-file cow paths. Pigs, however, do not possess the instinct to walk in single file (Grandin, 1994). Increased occurrence of highly excitable pigs is associated with both meat quality and animal welfare problems. Heavy muscled, excitable pigs will constantly balk in the restraining chute making it necessary for plant personnel to apply increasingly stressful handling practices in an effort to maintain constant delivery of pigs to the stunner at rates of (or above) 1,000 pigs per hour. Rough handling of pigs in the stunning chute during the last few minutes before exsanguination will increase the occurrence of PSE meat even in pigs resistant to the stress gene (Tarrant, 1989). Use of two stunning systems which operate at 500 pigs per hour will greatly reduce problems associated with meat quality and animal welfare. Appropriate design of restraining chutes (a side-by-side loading chute as described in *Pre-transport handling at the production site*) will assist proper delivery of pigs to the stunner.

Rendering the animal insensible

Under the Humane Methods of Slaughter Act of 1978, all animals slaughtered under federal inspection must be rendered insensible prior to exsanguination.

Electrical stunning is the most common method of stunning slaughter pigs in the United States. With electric stunning, sufficient amperage must pass through the pigs brain to induce an epileptic seizure resulting in instantaneous, painless, unconsciousness of the pig. Meat packers should use amperage, voltage, and frequency settings that will reliably induce insensibility. When electric stunning is done correctly, the animal will feel nothing. Insufficient amperage or poor current contact may paralyze the pig, however, the animal will not be insensible and will experience the feeling of a large electric shock and heart attack symptoms (Grandin, 1998e). To insure proper, humane stunning of pigs and to avoid potential meat quality problems, Grandin (1994) recommends use of a constant amperage power supply where the amperage is a constant 1.25 amps and the voltage varies with pig resistance. A minimum of 300 volts should be used for larger pigs and slightly lower voltages for lighter hogs. Grandin (1994) notes that packing plants should never reduce stunning amperage (for example, 0.5 amps) in an attempt to reduce blood splash because low amperages (or frequencies over 200 Hz) fail to induce instantaneous unconsciousness (Hoenderken, 1983).

Stunning and meat quality

A review of the effect of stunning on carcass and meat quality is provided by Gregory (1987). Carcass and meat quality defects influenced by stunning include bone fractures, blood splash, bruising, inadequate bleeding, and PSE meat. Stunning pigs with high voltages may result in broken shoulder blades if the animal is not supported off the floor. Hemorrhage associated with broken bones will require trimming, resulting in more time on the kill floor (at an elevated temper-

ature) which is detrimental to pork quality. Shattered scapulas can be avoided either by elevating the animal from the floor prior to stunning (via a restraining conveyor) or by reducing the length of time current is applied to the animal (2 seconds at 320 volts; Braathen and Johansen, 1984). High stunning currents administered to restrained pigs have also resulted in fractured thoracic vertebra and pelvis (Gregory, 1987).

Blood splash (ecchymosis) occurs when the physiological stress of stunning induces capillary rupture resulting in the appearance of small spots or large clots of blood in the lean tissue. High voltage stunning in a restraining conveyor and captive bolt stunning lead to intense muscular contraction and a high incidence of blood splash (Burson *et al.*, 1983; Larsen, 1983). A study by van der Wal (1978) reported that stimuli from the adrenal medulla continued after captive bolt stunning, increasing blood catecholamine concentrations and violent muscle contractions. These actions increase lactate concentrations in the blood, cause capillary hemorrhage (blood splash), and (or) carcass bruising. Gregory (1987) summarized that high voltage head to back stunning (240V at 50Hz) gave a reduction in the severity of blood splash compared to low voltage, high frequency (97V at 1700Hz) or low voltage, low frequency (80-125V at 50Hz) head only stunning. Larsen (1982) found less blood splash in carcasses from pigs stunned at 700V compared to those stunned at 300V. In these studies, the higher voltage could have been sufficient to induce cardiac arrest, stopping the heart altogether, eliminating the means (pump) to force blood through the ruptured blood vessels, thus generating a lower occurrence of blood

continued on page 5

continued from page 4

splash.

The operator of the stunning equipment must be attentive and well trained. The operator should keep the contacts clean and watch for worn cords and switches. The operator should avoid sliding the stunning wand over the animal during stun. Grandin (1994) refers to this as "double stunning" because it causes the pig to contract more than once. Double stunning can also occur if the electricity is turned on before the applicator is pressed firmly against the pig.

Reducing the amount of time between stunning and sticking minimizes the period when increased blood pressure provides more force for blood to leak through ruptured blood vessels (Burson *et al.* 1983). Many U.S. slaughter plants apply a horizontal stick before the pig is shackled as a means of reducing the stun to stick

interval. This will improve meat quality by reducing bruising and prevent struggling after stunning (Eilert, 1997). Grandin (1994) recommends bleeding pigs within 10 seconds after stunning (Burson *et al.*, 1983 advise 10 to 20 s) to minimize blood splash. Exsanguination as soon as possible after stunning is an important practice of humane slaughter, reducing the possibility an animal will regain consciousness during the initial stages of slaughter (Grandin, 1994). Wotton and Gregory (1986) found that after severing the brachiocephalic trunk and anterior vena cava, it took (on average) 18 seconds to induce a loss of brain function in 55 kg (avg. weight) pigs. Hoenderken (1983) reported that an experienced sticker could induce an isoelectric electrocorticogram in 12 to 20 seconds. A pig that is stuck incorrectly may take 55 to 60 seconds to lose responsiveness (Wotton and Gregory, 1986).

pH units/min while a rate of pH decline of 0.02 units/min (representing a marginal case of PSE) would increase the fraction of denatured myosin 7 fold. At a rate of pH fall of 0.1 units/min (severe PSE), this same extreme temperature abuse would only increase the fraction of denatured myosin a factor of 1.7. It is apparent that more rapid chilling has greater benefits for carcasses experiencing slower and intermediate pH declines.

Temperature and pH of a pork carcass may vary from the right to left side. According to van der Wal *et al.* (1995), initial temperature of the right *biceps femoris* and *semimembranosus* muscles were significantly higher than their counterparts on the left side yet *longissimus lumborum* and subcutaneous temperature remained equal. The pH of the right *semimembranosus* was also found to be significantly lower within the right ham. These differences in temperature and pH were maintained till at least 45 min *postmortem*. The authors discovered that the right and left differences in temperature and pH were a consequence of shackling the carcass on the right hind leg immediately following stunning. Care should be exercised in the design of pork quality experiments to include side randomization or to randomize right and left leg shackling.

Postmortem Temperature and pH

Classic work by Penny (1969) reported that *postmortem* rate of pH decline of PSE meat is approximately three times faster than normal, producing an intramuscular pH below six before the carcass temperature has time to cool below 37° C. The combination of low pH and high temperature results in muscle protein denaturation which leads to the reduced water holding capacity associated with PSE meat. The net electrical charge of myosin becomes minimal as the pH of meat nears the isoelectric point (pI) of myosin ($pI = 5.4$) resulting in a low water binding capacity (Wisner-Pedersen, 1987). Lean tissue possessing a low water holding capacity is detrimental to the meat processing industry as approximately 75% of all pork meat is sold in the processed form (Cannon *et al.*, 1995). Eilert (1997) suggested that an ideal plant would have the carcasses off of the slaughter floor within 20 to 25 min. The plant should do everything possible to reduce the length of time from stunning to chilling.

Offer (1991) generated a model for the formation of PSE meat reporting that the normal rate of pH fall in pork carcasses is about 0.01 pH units/min, corresponding to a rigor time of about 150 min (assuming a linear pH decline). A marginal case of

PSE (corresponding to a pH at 45 min \pm 6; Penny, 1969) would have a rate of pH decline of 0.02 units/min and an extreme case (achieving rigor in 15 min *postmortem*) would have a rate of decline of 0.1 pH units per min. Prerigor muscle is metabolically active generating heat from the conversion of glycogen to lactic acid and the hydrolysis of creatine phosphate and ATP (Essen-Gustavsson, 1993; Swatland, 1993). Prolonged contractile activity results in localized accumulation of heat, CO₂, and lactic acid which results in a higher than normal body temperature. At a very high rate of pH decline (> 0.067 pH units/min) the muscle temperature initially rises before falling as a consequence of this metabolic activity. Offer (1991) proposed that metabolic heat was proportional to the pH fall at a rate of 2°C/pH unit and found that more rapid chilling always decreased the amount of myosin denatured at rigor. Protein denaturation in a carcass with a relatively moderate pH decline and a prolonged period of temperature abuse may be greater than in a carcass experiencing more severe conditions (elevated temperature and rapid pH decline) for a shorter period duration. Offer (1991) reported that increasing the length of time to reach carcass *half-cooling* from 100 to 700 min produced an 18 fold increase in the fraction of denatured myosin when the rate of pH decline occurred at the normal rate of 0.01

Scalding of Carcasses

Tarrant (1993) concluded that factors affecting carcass temperature or delay chilling (such as ante-mortem handling, air temperature on the slaughter floor, and carcass scalding) require specific attention. In their study on the affect of scalding on subcutaneous and deep ham temperatures, van der Wal *et al.* (1993) found that scalding carcasses at 60°C for at least 5.5 to 7.5 minutes gave satisfactory dehairing results with the exception of carcasses of pigs slaughtered in autumn (during *hard hair* season). The latter required 9 min scalding to be dehaired sufficiently. Van der Wal (1993) reported that heat was absorbed superficially during scalding and can be removed quickly when scalding and scorching have finished. The group discovered that even an extremely long scalding time of 19 minutes only increased ham muscle temperature (5 cm below the skin in the *biceps femoris*) approximately 1°C.

continued on page 6

continued from page 5

The authors concluded that scalding pork carcasses for 9 minutes will not affect pork quality however, even a small increase of muscle temperature means that no heat was removed from the tissue over the time this increase was maintained.

Monin *et al.* (1995) tested differences in fresh pork quality as a result of pigs dehaired by scalding or singeing. During the process of singeing, the pigs body is passed through a series of gas burners before they are scraped and brushed (polished) in the usual way. The authors found no difference in 30 min post-mortem temperature or pH between the two dehairing methods. The singed carcasses showed slightly higher ultimate pH (20 h PM) in the *semimembranosus* and *aductor* and lower reflectance and higher water holding capacity in the *biceps femoris*. Monin *et al.* (1995) concluded that meat quality was slightly better in singed carcasses but differences in meat quality between the

dehairing methods was insignificant.

Eilert (1997) suggested that skinning of pork carcasses will improve meat quality in the sense that it will shorten the time prior to evisceration. Rapid emptying of the peritoneal cavity results in elimination of the visceral heat and more rapid carcass temperature decline and removal of the skin will result in more rapid transmittance of heat from the interior musculature. Troeger (1987) found the average temperature of skinned versus scalded carcasses was approximately 1°C cooler in the loin and ham at 30 min *postmortem*. At 1 h *postmortem*, the average pH of the *biceps femoris* of skinned carcasses was 6.23 and scalded carcasses 5.90, however, no difference was observed for ultimate pH. Skinned carcasses had a lower cooler shrink loss (1.8% versus 2.6%) compared to scalded carcasses. No differences were observed for 24 h light intensity of the *longissimus lumborum*, however after 4 days of vacuum packaging, the loins from skinned carcasses were significantly darker.

of purge and cooking loss without negatively affecting Warner-Bratzler shear force (WBS). The blast chilling process did not induce cold toughening and (or) decrease tenderness even though the carcasses used in this study were very lean possessing minimal external fat (10th rib fat depth = 1.34 cm).

Van der Wal *et al.* (1995) tested 3 different chilling regimes. A moderate conventional chilling regime (0-4°C; air velocity 0.5 m/s) served as a control and two forced chilling regimes; moderate rapid chilling (-5°C) for 2h or ultra rapid chilling (-30°C) for 30 min with a downward air velocity of 1, 2, or 4 m/s. After forced chilling, the carcasses were delivered to the cooler for equalization chilling (0-4°C, 0.5 m/s) till 24 h *postmortem*. The authors found a significant effect of forced chilling on muscle temperature compared to control carcass halves. The internal muscle temperatures after prolonged (120 min) moderate (-5°C) chilling were lower compared to ultra rapid (30 min) chilling (-30°C). An increase in air velocity resulted in a reduction of *longissimus lumborum* and *semimembranosus* temperature but not *biceps femoris* (believed to be due to increased external fat covering this muscle). Ultra rapid chilling generated less carcass shrink compared to conventional and moderate forced chilling, however, meat quality parameters (with the acceptance of variables related to tenderness) were not affected by chilling regime. Taylor *et al.* (1995) found no difference in quality associated with different cooling rates. Jones *et al.* (1993) reported that carcasses blast chilled for 3 h (-20 or -40°C with an air velocity of 5m/s) had *longissimus thoracis* muscles that were darker, had a higher protein solubility, less drip loss, shorter sarcomere lengths, and higher Warner-Bratzler shear force values compared to conventional chilling (1°C, air velocity of 1 m/s and forty, 60 s spray chill cycles set at 15 min intervals). Furthermore, muscle color was darker in the *longissimus thoracis* and *semimembranosus* muscles of carcasses blast chilled for 3 h at -40°C compared to the conventional spray-chill treatment. The authors note that these color differences would be "on the borderline of detection" using subjective visual appraisal. In another study, Jeremiah *et al.* (1992) reported that carcass sides blast chilled for 1h (-20 or -40°C)

Carcass Chilling

Grandin (1994) believes that handling practices account for 10 to 15% of the variation in PSE meat and chilling practices account for 20 to 40%. Blast chilling of carcasses with forced air at temperatures well below freezing has been used as a means of rapidly reducing intramuscular temperature to reduce the occurrence of PSE meat. Eilert (1997) stated that more rapid chilling procedures will not prevent PSE because, in severe cases of PSE, the damage has already occurred before the carcasses enter the cooler. As stated in the previous section, more rapid chilling has greater benefits for improving quality in carcasses experiencing slow and intermediate pH declines.

Rapid chilling of pork muscle may generate a toughening effect through muscle shortening (cold shortening). Jones *et al.* (1993) found no significant differences in mean Warner-Bratzler shear force values for blast chilled versus conventional chilled carcasses. However, blast chill treatments did increase the variability in shear values. The standard errors for shear values of blast chilled carcasses were 2 - 3 times greater than those of carcasses spray chilled at conventional chilling rates.

Jones *et al.* (1991) submerged pork

sides in liquid nitrogen at 40 min *postmortem* as an extreme method of removing carcass heat. Carcass sides were completely submerged for 1 - 3 min. The group found no significant treatment differences in meat color, protein solubility, or sarcomere length. Warner-Bratzler shear force was significantly increased in chops obtained from sides chilled by liquid nitrogen for one minute but surprisingly, no significant difference was noted between conventional chill and 3 min submersion in liquid nitrogen. A second study revealed that liquid nitrogen chilling had no effect on meat color, drip loss, protein solubility, sarcomere length, or shear value. Jones *et al.* (1993) noted that blast chilling can induce cold shortening of the muscles if temperatures fall below 10°C when the muscle pH is above 6.0. Cold shortening may be more of problem when heavy muscled, lean carcasses are blast chilled. Less external fat will allow a more rapid release of carcass heat and a greater susceptibility to muscle cold shortening. McFarlane and Unruh (1996) reported blast chilled pork carcasses (1 hr at -25°C and then chilled 23 hr at 1°C) tended to have a higher 24-h pH than carcasses chilled in the conventional manner (24 h at 1°C). Increasing the chilling rate also reduced the combined percentage

continued on page 7

continued from page 6

were more tender and rated higher in overall palatability than sides blast-chilled for 2 or 3 h. The authors were quick to clarify that the magnitude of these differences had limited practical significance as the palatability differences between short and extreme (-40°C for 3 h) blast-chill times were inconsequential.

Accelerated carcass chilling will most often have little or no significant effect on final lactic acid concentrations in the muscle, however, it will reduce the rate of lactic acid formation (Wisner-Pedersen, 1987). Rapid chilling of pork carcasses will most always reduce carcass cooler shrink, however, the effects on pork quality parameters are limited and conflicting in the literature.

Conclusions

The term welfare is relevant only when an animal is alive but death during handling and transport is usually preceded by a period of poor welfare (Broom, 1993). Improving the welfare of market pigs will generate less stress on the pig and give less stress to the individuals handling these pigs. Handling of pigs greatly improved when the United States entered the Japanese export trade. Grandin (1993) declares that when slaughter plant managers watched a Japanese grader reject up to 40% of their pork loins due to PSE a strong economic incentive was created to improve handling. Simple changes in handling procedures such as showering and resting pigs enabled 10% more pork to be exported to Japan (Grandin, 1993). Pre-slaughter treatments are much easier to control than the *postmortem* biochemical reactions triggered by stressful handling (Honkavaara, 1989c). Attention to the critical points outlined in this review is necessary to produce a high quality food product and make pork the meat of choice in the next millennium.

Literature Cited

- Baxter, M. 1985. Social space requirements for pigs. P. 116-127. In: Social Space for domestic Animals. R. Zayan Edit. Martinus Nijhoff Publ. Boston, MA.
- Becker, B. A., H. F. Mayes, G. L. Hahn, J. A. Nienaber, G. W. Jesse, M. E. Anderson, H. Heymann, and H. B. Hedrick. 1989. Effect of fasting and transport on various physiological parameters and meat quality of slaughter hogs. *J. Anim. Sci.* 67:334-341.
- Braathen, O. S. and J. Johansen. 1984. The effect of short or long electrical stunning times upon pork quality. *Proc. 30th European Meeting of Meat Research Workers.* 30:22-23 (as cited by Gregory, 1987).
- Bradshaw, R. H., R. F. Parrot, J. A. Goode, D. M. Loyd, R. G. Rodway, and D. M. Broom. 1996. Effects of mixing and duration of journey on the welfare of pigs during transport. P. 95-100. In: *Proc. EU-Seminar: New information on welfare and meat quality of pigs as related to handling, transport and lairage conditions.* Braunschweig: Bundesforschungsanstalt für Landwirtschaft Braunschweig-Völkeroede.
- Broom, D. M. 1993. Welfare assessment and welfare problem areas during handling and transport. P. 35-42. In: *Livestock Handling and Transport.* Edit. T. Grandin. CAB International. Wallingford, UK.
- Burson, D. E., M. C. Hunt, D. E. Schafer, D. Beckwith, and J. R. Garrison. 1983. Effects of stunning method and time interval from stunning to exsanguination on blood splashing in pork. *J. Anim. Sci.* 57:918-921.
- Cannon, J. E., J. B. Morgan, J. Heavner, F. K. McKeith, G. C. Smith, and D. L. Meeker. 1995. Pork quality audit: A review of the factors influencing pork quality. *J. Muscle Foods.* 6:369-402.
- De Smet, S. M., H. Pauwels, S. De Bie, D. I. Deymeyer, J. Callewier, and W. Eeckhout. 1996. Effect of halothane genotype, breed, feed withdrawal, and lairage on pork quality of Belgian slaughter pigs. *J. Anim. Sci.* 74:1854-1863.
- Eilert, S. J. 1997. What quality controls are working in the plant? P. 59-63. In: *Proc Pork Quality Summit.* July 8-9. National Pork Producers Council. Des Moines, IA.
- Eikelenboom, G., A. H. Bolink, and W. Sybesma. 1991. Effects of feed withdrawal before delivery on pork quality and carcass yield. *Meat Sci.* 29:25-30.
- Elliott, R. J. and J. Patton. 1968. The effects of road transportation and lairage treatment on pig muscle. *Proc. European Meeting of Meat Research Workers.* 14:397-405.
- Essen-Gustavsson, B. 1993. Muscle-fiber characteristics in pigs and relationships to meat-quality parameters - Review. P. 140-159. In: *Pork Quality: Genetic and Metabolic Factors.* Edit. E. Puolanne, D. I. Demeyer, M. Ruusunen, and S. Ellis. CAB International.
- Enfalt, A. C., K. Lundstrom, and U. Engstrand. 1993. Early *post mortem* pH decrease in porcine *M. Longissimus dorsi* of PSE, normal and DFD quality. *Meat Sci.* 34:131-143.
- Grandin, T. 1989. Behavioral principles of livestock handling. In: *Professional Animal Scientist.* Dec. American Registry of Professional Animal Scientists.
- Grandin, T. 1993. Introduction: Management and economic factors of handling and transport. P. 1-9. In: *Livestock Handling and Transport.* Edit. T. Grandin. CAB International. Wallingford, UK.
- Grandin, T. 1994. Methods to reduce PSE and Bloodsplash. *Proc. Allen D. Leman Swine Confr.* University of MN. 21:206-209.
- Grandin, T. 1997. Assessment of stress during handling and transport. *J. Anim. Sci.* 75:249-257.
- Grandin, T. 1998a. Dealing with excitable pigs. Temple Grandin's web page. www.grandin.com/meat/pigs/excite.pig.1.html.
- Grandin, T. 1998b. Loading and unloading animals properly. Temple Grandin's web page. www.grandin.com/behaviour/unload.truck.html.
- Grandin, T. 1998c. Principles of low stress restraint for pigs. Temple Grandin's web page. www.grandin.com/meat/pigs/rest.princ.2.html.
- Grandin, T. 1998d. Recommended trucking practices. Temple Grandin's web page. www.grandin.com/behaviour/rec.truck.html.
- Grandin, T. 1998e. Electric stunning of pigs. Temple Grandin's web page. www.grandin.com/meat/pigs/elec.stun.1.html.
- Gregory, N. G. 1987. Effect of stunning on carcass and meat quality. P. 265-272. In: *Evaluation and control of meat quality in pigs.* Edit. P. V. Tarrant, G. Eikelenboom, and G. Monin. Martinus Nijhoff Publ. Boston, MA.
- Guise, H. J. and P. D. Warris. 1989. A note on the stocking density and temperature on meat quality in pigs. *Anim. Prod.* 48:480-482.
- Guise, H. J. and R. H. C. Penny. 1989a. Factors influencing the welfare and carcass and meat quality of pigs. 1. The effects of stocking density in transit and the use of electric goats. *Anim. Prod.* 49:511-515.
- Guise, H. J. and R. H. C. Penny. 1989b. Factors influencing the welfare and carcass and meat quality of pigs. 2. Mixing unfamiliar pigs. *Anim. Prod.* 48:480-482.
- Hemsworth, P. H. 1993. Behavioural principles of pig handling. P. 197 - 211. In: *Livestock Handling and Transport.* Edit. T. Grandin. CAB International. Wallingford, UK.
- Hoenderken, R. 1983. Electrical and carbon dioxide stunning of pigs for slaughter. P. 59-63. In: *Stunning of animals for slaughter.* Edit. G. Eikelenboom. Martinus Nijhoff Publ. Boston, MA.
- Honkavaara, M. 1989a. Influence of selection phase, fasting and transport on porcine stress and on the development of PSE pork. *J. Agric. Sci. Finland.* 61:415-423.
- Honkavaara, M. 1989b. Influence of lairage on blood composition of pig and on the development of PSE pork. *J. Agric. Sci. Finland.* 61:425-432.
- Honkavaara, M. 1989c. Influence of carcass temperature, glycogenolysis and glycolysis 45 min *postmortem* on the development of PSE pork. *J. Agric. Sci. Finland.* 61:433-440.
- Hunter, E. J., H. L. Riches, H. J. Guise, and R. H. C. Penny. 1997. The behaviour of pigs in lairage in relation to their post-weaning management: results of a postal survey. *Anim. Welfare.* 6:139-144.
- Jeremiah, L. E., S. D. M. Jones, G. Kruger, A. K. W. Tong, and R. Gibson. 1992. The effects of gender and blast-chilling time and temperature on cooking properties and palatability of pork *longissimus* muscle. *Can. J. Anim. Sci.* 72:501-506.
- Jones, S. D. M., R. E. Rempala, and C. R. Haworth. 1985. Effects of water restriction on carcass shrink and pork quality. *Can. J. Anim. Sci.* 65:613-618.
- Jones, S. D. M., G. G. Greer, L. E. Jeremiah, A. C. Murray, and W. M. Robertson. 1991. Cryogenic chilling of pork carcasses: Effects on muscle quality, bacterial populations and palatability. *Meat Sci.* 29:1-16.
- Jones, S. D. M., L. E. Jeremiah, and W. M. Robertson. 1993. The effects of spray and blast-chilling on carcass shrinkage and pork muscle quality. *Meat Sci.* 34:351-362.
- Kilgour, R. and C. Dalton. 1984. *Livestock Behaviour: a practical guide.* Granada Publishing Ltd. New York, NY.
- Klont, R. E. and E. Lambooy. 1995. Influence of preslaughter muscle temperature on muscle metabolism and meat quality in anesthetized pigs of different halothane genotypes. *J. Anim. Sci.* 73:96-107.
- Lambooy, E. and G. van Putten. 1993. Transport of pigs. P. 213-231. In: *Livestock Handling and Transport.* Edit. T. Grandin. CAB International. Wallingford, UK.
- Larson, H. K. 1983. Comparison of 300 volt stunning, 700 volt automatic stunning and CO₂ compact stunning, with respect to quality parameters, blood splashing, fractures and meat quality. P. 73-81. In: *Stunning of animals for slaughter.* Edit. G. Eikelenboom. Martinus Nijhoff Publ. Boston, MA.
- March, B. B., R. G. Cassens, R. G. Kauffman, and E. J. Briskey. 1972. Hot boning and pork tenderness. *J. Food Sci.* 37:179-180.



- Martoccia, L., G. Brambilla, A. Macri, G. Moccia, and E. Cosentino. 1995. The effect of transport on some metabolic parameters and meat quality in pigs. *Meat Sci.* 40:271-277.
- McFarlane, B. J. and J. A. Unruh. 1996. Effects of blast chilling and *postmortem* calcium chloride injection on tenderness of pork longissimus muscle. *J. Anim. Sci.* 74:1842-1845.
- Messe, G. B. and R. Ewbank. 1973. The establishment and nature of the dominance hierarchy in the domesticated pig. *Anim Behav.* 21:326-334.
- McGlone, J. J., J. L. Salak, E. A. Lumpkin, R. L. Nicholson, M. Gibson, and R. L. Norman. 1993. Shipping stress and social status effects on pig performance, plasma cortisol, natural killer cell activity and leukocyte numbers. *J. Anim. Sci.* 71:888-896.
- Mickwitz, von G. 1982. Various transport conditions and their influence on physiological reactions. P. 45-56. In: Transport of animals intended for breeding, production and slaughter. Edit. R. Moss. Martinus Hijhoff Publ. Boston, MA.
- Mitchell, G. and J. J. A. Heffron. 1982. Porcine stress syndromes, porcine stress syndrome, malignant hyperthermia, and pale, soft, exudative pork syndrome. *Adv Food Res.* 28:167-230.
- Monin, G. A. Talmant, P. Aillery, and G. Collas. 1995. Effects on carcass weight and meat quality of pigs dehaired by scalding or singeing *post-mortem*. *Meat Sci.* 39:247-254.
- Murray, A. C., S. D. M. Jones, and A. P. Sather. 1989. The effects of preslaughter feed restriction and genotype for stress susceptibility on pork lean quality and composition. *Can. J. Anim. Sci.* 69:83-91.
- Nanni Costa, L., D. P. Lo Fiego, A. De Grossi, and V. Russo. 1996. Effect of loading method, stocking density and temperature on carcass and meat quality in heavy pigs. P. 83-93. In: *Proc. EU-Seminar: New information on welfare and meat quality of pigs as related to handling, transport and lairage conditions.* Braunschweig: Bundesforschungsanstalt für Landwirtschaft Braunschweig-Volkenrode.
- Offer, G. 1991. Modeling the formation of pale, soft and exudative meat: Effects of chilling regime and rate and extent of glycolysis. *Meat Sci.* 30:157-184.
- Pearson, A. M. 1987. Muscle function and *postmortem* changes. P. 155-191. In: *The Science of Meat and Meat Products.* 3rd Edit. Price and Schwiebert Edit. Food and Nutrition Press Inc. Westport, CN.
- Penny, I. F. 1969. Protein denaturation and water holding capacity in pork muscle. *J. Food Technol.* 4:269-273.
- Schutte, A., A. Mergens, U. Pott, and S. Venthien. 1996. Effect of different kinds of showering in lairage on physiological and meat quality parameters, taking climatic circumstances into account. P. 181-205. In: *Proc. EU-Seminar: New information on welfare and meat quality of pigs as related to handling, transport and lairage conditions.* Braunschweig: Bundesforschungsanstalt für Landwirtschaft Braunschweig-Volkenrode.
- Stephens, D. B. and G. C. Perry. 1990. The effects of restraint, handling, simulated and real transport in the pig (with reference to man and other species). *Appl. Anim. Behav. Sci.* 28:41-55.
- Swatland, H. J. 1993. Growth physiology and *post-mortem* metabolism in porcine muscle. P. 115-139. In: *Pork Quality: Genetic and Metabolic Factors.* Edit. E. Puolanne, D. I. Demeyer, M. Ruusunen, and S. Ellis. CAB International.
- Tarrant, P. V. 1989. The effects of handling, transport, slaughter and chilling on meat quality and yield in pigs. P. 1-25. In: *Manipulating Pig Production II.* Proc. Biennial Conference of the Australian Pig Science Assoc. Edit. J. L. Barnett and D. P. Hennessy. Warrabee, Victoria, Australia.
- Tarrant, P. V. 1993. An overview of production, slaughter and processing factors that effect pork quality - General review. P. 1 - 21. In: *Pork Quality: Genetic and Metabolic Factors.* Edit. E. Puolanne, D. I. Demeyer, M. Ruusunen, and S. Ellis. CAB International.
- Taylor, A. A., G. R. Nute, and C. C. Warkup. 1995. The effect of chilling, electrical stimulation and conditioning on pork eating quality. *Meat Sci.* 39:339-347.
- Troeger, K. 1987. Skinning versus scalding: influence on pork quality. P. 111 - 126. In: *Accelerated Processing of Meat.* Edit. R. Romita, C. Valin, and A. A. Taylor. Elsevier Applied Science. New York, NY.
- van der Wal, P. G., G. van Beek, C. H. Veerkamp, and G. Wijngaards. 1993. The effect of scalding on subcutaneous and ham temperatures and ultimate pork quality. *Meat Sci.* 34:395-402.
- van der Wal, P. G. 1978. Chemical and physiological aspect of pig stunning in relation to meat quality - A review. *Meat Sci.* 2:19-30.
- van der Wal, P. G., B. Engel, G. van Beek, and C. H. Veercamp. 1995. Chilling pig carcasses: Effects on temperature, weight loss and ultimate meat quality. *Meat Sci.* 40:193-202.
- van Putten, G. 1982. Handling of slaughter pigs prior to loading and during loading on a lorry. P. 15-27. In: *Transport of animals intended for breeding, production and slaughter.* Edit. R. Moss. Martinus Hijhoff Publ. Boston, MA.
- Weeding, C. M., H. J. Guise, and R. H. C. Penny. 1993. Factors influencing the welfare and carcass and meat quality of pigs: the use of water sprays in lairage. *Anim. Prod.* 56:393-397.
- Warris, P. D., S. C. Kestin, and J. M. Robinson. 1983. A note on the influence of rearing environment on meat quality in pigs. *Meat Sci.* 9:271-279.
- Warris, P. D. 1987. The effect of time and conditions of transport and lairage on pig meat quality. P. 245-264. In: *Evaluation and control of meat quality in pigs.* Edit. P. V. Tarrant, G. Eikelenboom, and G. Monin. Martinus Nijhoff Publ. Boston, MA.
- Warris, P. D., S. N. Brown, E. A. Bevis, and S. C. Kestin. 1990. The influence of pre-slaughter transport and lairage on meat quality in pigs of two genotypes. *Anim. Prod.* 50:165-172.
- Warris, P. D., S. N. Brown, J. E. Edwards, and T. G. Knowles. 1996. Effect of lairage time on levels of stress and meat quality in pigs. P. 163-170. In: *Proc. EU-Seminar: New information on welfare and meat quality of pigs as related to handling, transport and lairage conditions.* Braunschweig: Bundesforschungsanstalt für Landwirtschaft Braunschweig-Volkenrode.
- Williams, J. M. 1985. Causes and effects of porcine stress syndromes. P.175-179. In *The Veterinary Annual.* 25th issue. Bristol, Scientifica (as cited by Tarrant, 1993).
- Wismer-Pedersen, J. 1987. Chemistry of animal tissue - Part 5. Water. P. 141-154. In: *The Science of Meat and Meat Products.* 3rd Ed. Edit. J. F. Price and B. S. Schweigert. Food and Nutrition Press, Inc. Westport, CN.
- Wotton, S. B. and N. G. Gregory. 1986. Pig slaughtering procedures: time to loss of brain responsiveness after exsanguination or cardiac arrest. *Res. Vet. Sci.* 40:148-151.

For more information contact:



National Pork
Producers Council
P.O. Box 10383
Des Moines, Iowa USA
515 223 2600
Fax: 515 223 2646
Internet: pork@nppc.org
Web: <http://www.nppc.org/>

© 1998 National Pork Producers Council

National Pork Producers Council programs are supported in cooperation with the National Pork Board.